

Table 10.2
Kaposi Sarcoma

Number of Cases

By SEER Location and Year of Diagnosis, 1975-2006

All Races, Males, Ages 20-54

Year of Diagnosis

SEER Location	1975-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001	2002-2003	2004-2006
SEER 9 ^b	13	26	254	806	1480	1994	2104	1921	1676	753	317	344	257	418
SEER 9 except SF-Oakland	11	5	35	207	378	694	773	909	809	409	196	225	182	286
San Francisco-Oakland SMSA ^c	<4	21	219	599	1102	1300	1331	1012	867	344	121	119	75	132
San Francisco City/County	<4	18	193	520	924	1041	1051	748	649	236	76	73	51	92

Age-Adjusted Cancer Incidence Rates^a

By SEER Location and Year of Diagnosis, 1975-2006

All Races, Males, Ages 20-54

Year of Diagnosis

SEER Location	1975-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001	2002-2003	2004-2006
SEER 9 ^b	0.1	0.3	2.4	7.4	13.1	16.9	17.1	14.9	12.7	5.6	2.3	2.5	1.9	2.0
SEER 9 except SF-Oakland	0.1	0.1	0.4	2.2	3.9	6.9	7.4	8.4	7.2	3.6	1.7	2.0	1.6	1.6
San Francisco-Oakland SMSA ^c	-	1.3	12.1	32.3	57.3	64.4	63.9	46.8	40.2	15.8	5.4	5.2	3.3	3.9
San Francisco City/County	-	5.3	47.8	125.1	214.0	239.1	240.3	165.3	142.7	51.5	16.1	15.4	10.5	12.6

^a Rates are per 100,000 males aged 20-54 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).^b The SEER 9 areas are San Francisco-Oakland SMSA, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.^c The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi Sarcoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.